



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

$BR.a=c^2$, $AP.a=AR.a=bc$, $BP.a=a^2-b^2$, $CR.a=a^2-c^2$, i. e., $PR.a=2bccosA$; (4) hence BA touches circle ARC , which contains a Brocard-point of ABC ; similarly for CA and circle APB ; (5) $BR.CR'$, $AR''=abc=CP.BP'$, AP'' (where R' , R'' , P' , P'' correspond to R , P , on CA , AB , respectively); K , K' are the Brocard constants ($k=a^2+b^2+c^2$) of ABC , $A'B'C'$: then $K'-K=\Delta^2/R^2$.

181. Proposed by WILLIAM HOOVER, A. M., Ph. D., Professor of Mathematics and Astronomy, Ohio University, Athens, Ohio.

Prove that the extremities of the latera recta of all ellipses having a given major axis $2a$ lie on the parabola $x^2=-a(y-a)$.

CALCULUS.

142. Proposed by J. SCHEFFER, A. M., Hagerstown, Md.

Solve the differential equation,

$$(a-x)\frac{dz}{dx} + (b-y)\frac{dz}{dy} = c-z.$$

143. Proposed by L. C. WALKER, A. M., Petaluma High School, Petaluma, Cal.

Find the area of greatest ellipse that can be inscribed in a given semicircle.

144. Proposed by G. B. M. ZERR, A. M., Ph. D., Professor of Chemistry and Physics, The Temple College, Philadelphia, Pa.

Find the volume of the sphere, $x^2+y^2+z^2=2az$, (a) within the paraboloid $z=Ax^2+By^2$; (b) within the cone $z^2=Ax^2+By^2$.

MECHANICS.

134. Proposed by G. B. M. ZERR, A. M., Ph. D., Professor of Chemistry and Physics, The Temple College, Philadelphia, Pa.

If $pv=Rt-b/tv$ be the equation for CO_2 gas, find the total, external and internal work done in compressing the gas from 102 to 136 atmospheres at a constant temperature $17^\circ C$ and constant volume, $R=63.23$, $b=481600$ for CO_2 .

135. Proposed by F. P. MATZ, Sc. D., Ph. D., Professor of Mathematics and Astronomy in Defiance College Defiance, Ohio.

What force acting at an inclination ω with a horizontal line on the center of a wheel of given weight will roll the wheel over an immovable cylindric log whose diameter is $(1/m)$ th that of the wheel?

DIOPHANTINE ANALYSIS.

91. Proposed by F. P. MATZ, Sc. D., Ph. D., Professor of Mathematics and Astronomy in Defiance College, Defiance, Ohio.

There are two unequal square numbers the sum of whose sum, difference, product, and quotient, is a square. Find the two numbers.